Patrick Moore, MD, FACS

Breast Surgery Post-Operative Instructions (With a Drain)

PAIN MEDICATION

- You will be given a prescription for pain medication as needed. Drink a lot of water to help prevent constipation. You may take laxatives and/or stool softeners as needed.
- 2. When the pain decreases, you can change to something like Extra Strength Tylenol for pain relief.
- 3. If you experience nausea and/or vomiting, maintain your diet on clear liquids and call the office if the nausea and vomiting become persistent.

DRIVING

- 1. Do NOT drive.
- 2. Do NOT drive while taking the prescription pain medication.

DRESSINGS

1. You may take a sponge bath to clean around the dressing and drain area.

DRAIN

- 1. Record the Date, Time and Output each time the drain is emptied.
- 2. Call the office when the output amount is *less than 30 ml* over a 24 hour period to have the drain removed.
- 3. Change the dressing around the drain daily, if instructed to do so.

ACTIVITY

Keep activity light. You may walk outside or upstairs. No lifting over 10 lbs.

DIET

Start with clear liquids and add in solid food as tolerated.

Patrick Moore, MD, FACS

Breast Surgery Postoperative Instructions (With a Drain)

PAGE THE DOCTOR (

IF YOU EXPERIENCE:

- 1. Unusual color and/or discharge from the incision.
- 2. A fever over 101 degrees and chills.
- 3. A dramatic increase in your pain.
- 4. Persistent nausea and vomiting.
- 5. Please call us to schedule a follow- up appointment.

Call us to schedule a follow-up appointment within 2 weeks following your discharge.

Please call Dr. Moore's pager number if you have ANY serious concerns or guestions after surgery.